



Early Journal Content on JSTOR, Free to Anyone in the World

This article is one of nearly 500,000 scholarly works digitized and made freely available to everyone in the world by JSTOR.

Known as the Early Journal Content, this set of works include research articles, news, letters, and other writings published in more than 200 of the oldest leading academic journals. The works date from the mid-seventeenth to the early twentieth centuries.

We encourage people to read and share the Early Journal Content openly and to tell others that this resource exists. People may post this content online or redistribute in any way for non-commercial purposes.

Read more about Early Journal Content at <http://about.jstor.org/participate-jstor/individuals/early-journal-content>.

JSTOR is a digital library of academic journals, books, and primary source objects. JSTOR helps people discover, use, and build upon a wide range of content through a powerful research and teaching platform, and preserves this content for future generations. JSTOR is part of ITHAKA, a not-for-profit organization that also includes Ithaka S+R and Portico. For more information about JSTOR, please contact support@jstor.org.

$$= \frac{2\pi}{n} \sqrt{4r^2 \sin^2 \frac{\pi}{n} + \frac{d^2}{n^2}} \frac{mn(mn-1)}{2} = \pi m(mn-1) \sqrt{4r^2 \sin^2 \frac{\pi}{n} + \frac{d^2}{n^2}}$$

$$= \pi m^2 \sqrt{4\pi^2 r^4 + d^2}, \text{ when } n \text{ is infinite but } r=1 \text{ and } m=20, d=1.$$

$\therefore S=400\pi/(4\pi^2+1)=7995.12$ feet approximately.

[The above revised solution of Professor Zerr's now agrees in result with that of Professor Hume previously published.—Editor.]

17. Proposed by H. W. DRAUΘHON, Clinton, Louisiana

Find the volume generated by revolving a circular segment whose base is a given chord (2a), about any diameter as an axis.

II. First Solution by F. P. MATZ, M. Sc., Ph. D., Professor of Mathematics and Astronomy, New Windsor College, New Windsor, Maryland.

Let $\angle ECH=A$, $CA=r$, $AE=a$, $CE=x$, and $\angle ACD=\theta$; then $x'=r \cos \theta = \sqrt{r^2-a^2}$, and the required volume generated becomes

$$V=2(2\pi x \sin A) \int_x^r \sqrt{r^2-x^2} dx = 4\pi \sin A \left[-\frac{1}{2} (r^2-x^2)^{\frac{3}{2}} \right]_x^r = \frac{4}{3}\pi a^3 \sin A.$$

SECOND SOLUTION.

Put $\angle CAE=\phi$; then $x=r \sin \phi$, $dx=r \cos \phi d\phi$, and $\sqrt{r^2-x^2}=r \cos \phi$. The limits of integration are $\frac{1}{2}\pi$ and $\cos^{-1}(a/r)=\phi'$; and after obvious transformations, we have

$$V=4\pi \sin A \int_{\phi'}^{\frac{1}{2}\pi} r^3 \cos^2 \phi \sin \phi d\phi = 4\pi \sin A \left[-\frac{1}{3} r^3 \cos^3 \phi \right]_{\phi'}^{\frac{1}{2}\pi} = \frac{4}{3}\pi a^3 \sin A.$$

THIRD SOLUTION.

Represent the center of gravity of the segment $ADBE$ by G ; then from Mechanics, we have $CG = \frac{\frac{1}{2}\pi(2a)^3}{r^2 \sin^{-1}(a/r) - a\sqrt{r^2-a^2}}$, in which the denominator is the area of the segment $ADBE$, $=A$. Since $FG=CG \sin A = (2a^3/3A) \sin A$, the *Centrobaryc Method* gives for the volume generated $V=2\pi(2a^3/3A) \sin A \times A = \frac{4}{3}\pi a^3 \sin A$.

[In the solution by the Proposer, previously published the lower integral should have been $\sqrt{r^2-a^2}$. There was a misprint.—Editor.]

Note on Prob. No. 13 by JOHN DOLMAN, Jr., Philadelphia, Pennsylvania.

"It may be worth noting that the solution of No. 13 is incomplete in failing to state that the value of x found makes y a minimum only when angle EAD equals or is less than $\cos^{-1} \frac{4}{5}$ [considering the angle negative when D is west of E]. In all other cases the value, $x=\sin^{-1} \frac{4}{5}$, makes y a maximum and the steamers could meet in two possible points. All this, together with the critical value found, may be seen from inspection without any analytical work."

PROBLEMS.

27. Proposed by G. B. M. ZERR, A. M., Principal of Schools, Staunton, Virginia.

A runs around the circumference of a circular field with velocity m feet; B starts from the centre with velocity $n > m$ feet to catch A . The straight line joining their positions always passes through the centre. Find the equation to the curve described by B , the distance he runs and the time occupied.